

DTP/94-64
hep-ph/9407387

QCD INTERCONNECTION EFFECTS IN HADRONIC W^+W^- AND $t\bar{t}$ EVENTS

VALERY A. KHOZE¹

Department of Physics, University of Durham, Durham DH1 3LE, England

Abstract

In the events of the type $e^+e^- \rightarrow W^+W^- \rightarrow 4 \text{ jets}$, $e^+e^- \rightarrow t\bar{t} \rightarrow bW^+\bar{b}W^-$, particle production could depend in a non-trivial way on the kinematics of the process. It is shown that QCD interference effects are negligible for energetic perturbative emission, but soft perturbative gluons and non-perturbative fragmentation could induce colour correlations. Possible consequences for LEP 2 and NLC events are briefly addressed.

¹Presented at the Int. Conference “Quark Confinement and the Hadron Spectrum”, Villa Olmo-Como, Italy, June 20-24, 1994.

In high-energy physics, “tomorrow belongs” to the detailed study of heavy unstable particles (W bosons, top quarks, SUSY particles, ...). An important aim of future experiments is the precise determination of their parameters, primarily masses. This requires a detailed understanding of production and decay mechanisms (including interference effects) and, in particular, of the effects arising from the large width of many of these objects, $\Gamma \sim O(1 \text{ GeV})$. This talk is concerned with the QCD interconnection phenomena that may occur when two unstable particles decay and hadronize close to each other in space and time. The word ‘interconnection’ is here introduced to cover those aspects of final-state particle production that are not dictated by the separate decays of unstable objects, but can only be understood in terms of the joint action of the two. (For further details and a long list of references see Refs. [1,2].)

Let us start from hadronic W^+W^- events. QCD interferences between W^+ and W^- undermine the traditional meaning of a W mass in the process $e^+e^- \rightarrow W^+W^- \rightarrow q_1\bar{q}_2q_3\bar{q}_4$. Specifically, it is not possible to subdivide the final state into two groups of particles, one of which is produced by the $q_1\bar{q}_2$ system of the W^+ decay and the other by the $q_3\bar{q}_4$ system of the W^- decay: some particles originate from the collective action of the two systems. Since a determination of the W mass is one of the main objectives of LEP 2, it is important to understand how large the ambiguities can be. A statistical error of 55 MeV per experiment is expected, so the precision of the theoretical predictions should match or exceed this accuracy. A complete description of interference effects is not possible since non-perturbative QCD is not well understood. The concept of colour interconnection/rearrangement is therefore useful to quantify effects. In a rearrangement two original colour singlets (such as $q_1\bar{q}_2$ and $q_3\bar{q}_4$) are transmuted into two new ones (such as $q_1\bar{q}_4$ and $q_3\bar{q}_2$). Subsequently each singlet system is assumed to hadronize independently according to the standard algorithms. Depending on whether a reconnection has occurred or not, the hadronic final state is then going to be somewhat different. For a detailed understanding of QCD interconnection effects one needs to examine the space-time picture of the process. It was shown in [1] that interference is negligibly small for energetic perturbative gluon emission. Firstly, the W^+ and W^- decay at separate times after production, which leads to large relative phases for radiation off the two constituents of a rearranged system, and a corresponding dampening of the QCD cascades. Secondly, within the perturbative scenario the colour transmutation appears only in order α_s^2 and is colour-suppressed. It was concluded that only a few low-energy particles could be affected. In order to understand the interconnection effects occurring at the hadronization stage, the standard Lund fragmentation model has been considerably extended and several alternative models for the space-time structure of the fragmentation process have been developed^[1]. Comparing different models with the no-reconnection scenario, it turns out that interconnection effects are very small. The change in the averaged charged multiplicity is at the level of a per cent or less, and similar statements hold for rapidity distributions, thrust distributions and so on. The total contribution to the systematic error on the W mass reconstruction may be

as large as 40 MeV. This is good news. Otherwise, LEP 2 would not have significant advantages in the measurements of M_W over hadronic machines where the accuracy is steadily improving (the current combined results give $M_W = 80.23 \pm 0.18$ GeV and with the increase in statistics at the Tevatron further improvements are expected).

Clearly, colour rearrangement effects are interesting in their own right, for instance, as a new probe of the non-perturbative QCD dynamics. However, the standard measures considered in [1] seem to be below the experimental precision one may expect at LEP 2. A more optimistic conclusion has been reached in Ref. [3], where some specific ways to disentangle colour reconnection phenomena were proposed. But personally I still believe that one will need good luck in order to establish the nature and size of the QCD rearrangement effects in real-life experiments.

We turn now to the process $e^+e^- \rightarrow t\bar{t} \rightarrow bW^+\bar{b}W^-$ [2]. We assume that the W 's decay leptonically, so the colour flow is generated only by the t and b quarks. Further, we restrict ourselves to the region a few GeV above the $t\bar{t}$ threshold. The interplay of several particle production sources is reminiscent of the colour rearrangement effects we have studied for $e^+e^- \rightarrow W^+W^-$ but there are important differences. From the onset, W^+W^- events consist of two separate colour singlets, $q_1\bar{q}_2$ and $q_3\bar{q}_4$, so that there is no logical imperative of an interplay between the two. Something extra has to happen to induce a colour rearrangement to $q_1\bar{q}_4$ and $q_3\bar{q}_2$ singlets, such as a perturbative exchange of gluons or a non-perturbative string overlap. This introduces a sizeable dependence on the space-time picture, i.e. on how far separated the W^+ and W^- decay vertices are. Except in the unlikely case that top-flavoured hadrons would have time to form, the process $e^+e^- \rightarrow t\bar{t} \rightarrow bW^+\bar{b}W^-$ only involves one colour singlet. Therefore an interplay is here inevitable, while a colour rearrangement of the above kind is impossible.

One of the main objectives of a Next Linear e^+e^- Collider will be to determine the m_t with an accuracy of about 300 MeV. One method is to reconstruct the top invariant mass event by event, another is to measure the top momentum distribution. In either case, the colour flow restructuring could introduce the potentiality for a systematic bias in the m_t determination. In Ref. [2] we concentrated on the possible manifestations of the QCD interconnection effects in the distribution of the particle flow in the final state. As a specific example, we studied the multiplicity of leptonic top decays as a function of the angle between the b and $\bar{b}$ jets. The main conclusions of our study are:

- The interconnection phenomena should be readily visible in the variation of the average multiplicity as a function of the angle between the b and $\bar{b}$.
- A more detailed test is obtained by splitting the particle content in momentum bins. The high-momentum particles are mainly associated with the $\hat{t}\hat{b}$ and $\hat{t}\hat{\bar{b}}$ dipoles and follow the b and $\bar{b}$ directions, while the low-momentum ones are sensitive to the influence of the $\hat{b}\hat{\bar{b}}$ dipole.

- A correct description of the event shapes in top decay, combined with sensible reconstruction algorithms, should give errors on the top mass that are significantly less than 100 MeV.

The possibility of interference interconnection effects in the $t\bar{t}$ production is surely not restricted to the events studied here. One could discuss also hadronic W decays and/or the interferences with beam jets in $pp/p\bar{p} \rightarrow t\bar{t}$ events. The problem with these processes is that there are too many other uncertainties. At the moment, the main uncertainties come from the modelling of the non-perturbative fragmentation. However, the main conclusion that the QCD interconnection does not induce any sizeable restructuring of the final particle flows should remain valid for all of these cases.

Acknowledgements

I wish to thank Yu.L. Dokshitzer, V.S. Fadin, T. Sjöstrand and W.J. Stirling for enjoyable collaboration on aspects of the work presented here. This work was supported by the United Kingdom Particle Physics and Astronomy Research Council.

References

1. T. Sjöstrand and V.A. Khoze, *Z. Phys.* **C62** (1994) 281; *Phys. Rev. Lett.* **72** (1994) 28.
2. V.A. Khoze and T. Sjöstrand, *Phys. Lett.* **B328** (1994) 466.
3. G. Gustafson and J. Häkkinen, Lund preprint, LU TP 94-9.